

## Lunar Reconnaissance Orbiter (LRO) Observations with the Lunar Exploration Neutron Detector (LEND): Neutron Suppression Regions (NSR) and Polar Hydrogen

G. Chin<sup>1</sup>, I. G. Mitrofanov<sup>2</sup>, W. V. Boynton<sup>3</sup>, G. Droege<sup>3</sup>, J. B. Garvin<sup>1</sup>, D. V. Golovin<sup>2</sup>, L. G. Evans<sup>4</sup>, K. Harshman<sup>3</sup>, A. S. Kozyrev<sup>3</sup>, M. L. Litvak<sup>2</sup>, T. McClanahan<sup>1</sup>, G.M. Milikh<sup>5</sup>, R. Sagdeev<sup>5</sup>, A. B. Sanin<sup>2</sup>, V. Shevchenko<sup>7</sup>, V. Shvetsov<sup>8</sup>, D. Smith<sup>9</sup>, R. Starr<sup>6</sup>, J. Trombka<sup>1,5</sup> and M.Zuber<sup>9</sup>

<sup>1</sup>NASA Goddard Space Flight Center, Greenbelt, MD USA, [gordon.chin@nasa.gov](mailto:gordon.chin@nasa.gov), <sup>2</sup>Institute for Space Research, 117997 Moscow, Russia, <sup>3</sup>University of Arizona, Tucson, AZ USA, <sup>4</sup>Computer Science Corporation, Greenbelt, MD USA, <sup>5</sup>University of Maryland, College Park, MD USA, <sup>6</sup>Catholic University, Washington DC, USA, <sup>7</sup>Sternberg Astronomical Institute, Moscow, Russia, <sup>8</sup>Joint Institute of Nuclear Research, Dubna, Russia, <sup>9</sup>Massachusetts Institute of Technology, Cambridge, USA.


**Introduction:** Orbital detection of neutrons has become the dominant remote sensing technique for detecting and inferring H concentrations and its spatial distribution beneath planetary surfaces [Lawrence et al. (2010) *Icarus*, 205, pp. 195-209, Mitrofanov et al. (2007) *Science* 297(5578), 78-81].

Indications for the presence of localized and relatively high water content was provided by LRO and LCROSS. LEND identified *Cabeus*, as the most promising LCROSS impact site [Mitrofanov I. et al. (2010) *Science*, 330, 483], and instruments onboard LRO and LCROSS have measured signatures of water, H<sub>2</sub> and other volatiles in the impact plume [Colaprete A. et al. (2010) *Science*, 339, 463, Gladstone R. et al. (2010) *Science*, 330, 472].

**Neutron Suppression Regions (NSRs), as potential locations of polar Hydrogen.** Currently available LEND neutron data has identified localized *Neutron Suppression Regions*, or NSRs. The *Cabeus* NSR has a total area of about 700 km<sup>2</sup> (see Figure 1, *top*). Another well-observed NSR with an area about 1500 km<sup>2</sup> was detected within the polar crater known as *Shoemaker* (*middle*). The third example (*bottom*) of a well-defined NSR is the northern polar crater *Rozhdestvensky*.

The LEND data do not support a model that suggests PSRs are the sole traps of water molecules from the lunar exosphere. Paige et al. (2010) [*Science*, 330, 479] has shown that a layer of water-rich permafrost may exist below a thin upper-most layer of dry regolith even under conditions of solar irradiation. Also, based on analysis between LOLA and LEND, poleward-facing slopes on the Moon display a predominantly stronger neutron suppression in comparison with equator-facing slopes [McClanahan T. et al. (2011) Abstract #1970 of 42th LPSC].

ly. Surface relief is shown by gray shadows derived from LOLA altimetry.



**Figure 1:** LEND maps of epithermal neutrons for *Cabeus* (*top*), *Shoemaker* (*middle*) and at *Rozhdestvensky* (*bottom*) craters. Shades of blue corresponds to epithermal neutron flux. The boundaries of NSRs and PSRs are shown by red and white contours, respective-

